

# KINETICS AND MECHANISM OF THE OXIDATION OF THIOCYANATE ION BY DI- $\mu$ -OXO-TETRAKIS (1,10-PHENANTHROLINE)-DIMANGANESE (III, IV) ION IN ACID MEDIUM

Y. N. LOHDIP and J. F. IYUN

(Received 14 March 2002, Revision accepted 28 July 2002)

## ABSTRACT

The kinetics of oxidation of thiocyanate ion by di- $\mu$ -oxo-tetrakis(1,10-phenanthroline)dimanganese(III,IV) perchlorate in acid medium has been investigated. The reaction follows first order in both the oxidant and the reductant. The reaction is catalysed by hydrogen ion and the rate dependence given as  $k_p = a + b[H^+]$ . The reaction rate is not affected by changes in the ionic strength and dielectric constant of the reaction medium. The reaction did not induce polymerization of acrylamide and the presence of  $NO_3^-$  or  $HCOO^-$  had no effect on the rate. A plausible mechanism involving proton coupled electron transfer (PCET) is proposed

**Key words:** Kinetics, mechanism, oxidation, thiocyanate, di- $\mu$ -oxo-tetrakis(1,10-phenanthroline)-dimanganese (III,IV)

## INTRODUCTION

Interest in the chemistry of multinuclear manganese complexes has continued to increase (Reddy et. al., 1994; Arulsamy et. al., 1994; Jeffery et. al., 1994; Miao et. al., 1995; Goodson et. al., 1992; Vincent et. al., 1993; Dave and Czernuzewicz, 1994; Tanase and Lippard, 1995; Lohdip and Iyun, 1998, 2001). This interest is due to their relevance to redox activities in biosystems, usefulness in catalysis and molecular electronics (Babcock, 1987). Recently, we have investigated the redox reactions of the mixed valence complexes  $[L_2MnO_2MnL_2]^{3+}_{(aq)}$  (where L = 2,2'-bipyridyl or 1,10-phenanthroline) with hydroxy acids and thiols (Iyun et. al., 1997),  $H_2C_2O_4$  (Lohdip et. al., 1998), methionine (Lohdip and Iyun, 1998) and 2-mercaptobenzoic acid (Lohdip and Iyun, 2001). In continuation of our effort in this direction, we now report the results of the oxidation of SCN<sup>-</sup> by aqueous di- $\mu$ -oxo-tetrakis(1,10-phenanthroline)-dimanganese (III,IV) perchlorate. (referred to in the text as  $Mn^{III}O_2Mn^{IV}$ )

## EXPERIMENTAL

### Materials and reagents

The materials were of analytical or reagent grades and were used as received. The complex  $[(phen)_2Mn^{III}O_2Mn^{IV}(phen)_2](ClO_4)_3$  was prepared and characterized as described by Cooper and Calvin (1977). Sodium thiocyanate was standardized gravimetrically by oxidising it with bromine water and precipitated as barium sulphate (Bassett et. al., 1983). HCl was used to investigate the effect of hydrogen ions on the rate of the reaction while NaCl was employed to maintain the ionic strength constant at  $0.50 \text{ mol dm}^{-3}$ . All other chemicals were used as supplied

### Stoichiometric studies

The stoichiometry of the reaction was determined by spectrophotometric titration using the mole ratio method. Solutions containing known concentration of  $Mn^{III}O_2Mn^{IV}$  and varying concentrations of SCN<sup>-</sup> ( $0.55 - 3.86 \cdot 10^{-3} \text{ mol dm}^{-3}$ ) were reacted at  $[H^+] = 0.05 \text{ mol dm}^{-3}$  and  $I = 0.50 \text{ mol dm}^{-3}$  (NaCl). The absorbances were measured at 555 nm (on a solid state photometer, Trojanowicz et. al., 1988; Lohdip et. al., 1997), after completion of the reaction. The stoichiometry was evaluated from a plot of absorbance versus  $[SCN^-]/[Mn^{III}O_2Mn^{IV}]$ .

**Table 1:** Second order rate constants for the oxidation of thiocyanate ion by  $Mn^{III}O_2Mn^{IV}$ .  
 $[Mn^{III}O_2Mn^{IV}] = 4.403 \times 10^{-4} \text{ mol dm}^{-3}$ ;  $\lambda = 555 \text{ nm}$ ,  $T = 29.0 \pm 0.1^\circ\text{C}$ .

$10^3 [SCN^-]$ , $\text{mol dm}^{-3}$	$10^3 [H^+]$ , $\text{mol dm}^{-3}$	$I$ , $\text{mol dm}^{-3}$ (NaCl)	$10 k_2$ , $\text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$
2.76	5.0	0.5	1.10
4.14	5.0	0.5	1.11
5.51	5.0	0.5	1.11
6.89	5.0	0.5	1.18
8.27	5.0	0.5	1.19
9.65	5.0	0.5	1.21
10.10	5.0	0.5	1.21
12.41	5.0	0.5	1.21
13.78	5.0	0.5	1.22
16.54	5.0	0.5	1.23
19.30	5.0	0.5	1.23
22.05	5.0	0.5	1.21
10.10	1.0	0.5	0.81
10.10	2.0	0.5	0.89
10.10	7.0	0.5	1.36
10.10	10.0	0.5	2.05
10.10	12.0	0.5	2.27
10.10	15.0	0.5	2.63
10.10	5.0	0.1	1.16
10.10	5.0	0.2	1.20
10.10	5.0	0.7	1.18
10.10	5.0	1.0	1.17
10.10	5.0	1.2	1.23
10.10	5.0	1.5	1.21


### Kinetic studies

All kinetic runs were performed under pseudo-first-order conditions with the concentration of  $SCN^-$  in at least 10 fold excess over that of  $Mn^{III}O_2Mn^{IV}$ . The rate of reaction was monitored measuring the decrease in absorbance at 555 nm with a solid state photometer (Trojanowicz et. al., 1988; Lohdip et. al., 1997). Pseudo-first-order rate constants were obtained from logarithmic plots of absorbance differences against time. The temperature was maintained constant at  $29.0 \pm 0.1^\circ\text{C}$  with  $[H^+] = 0.05 \text{ mol dm}^{-3}$  and  $I = 0.5 \text{ mol dm}^{-3}$  (NaCl) unless otherwise stated.

## RESULTS AND DISCUSSION

### Stoichiometry

The results of the spectrophotometric titration indicated a 1 : 3 stoichiometry and the overall reaction is given by equation (1).



This stoichiometry is different from the one reported for most  $SCN^-$ /substrate reactions and for various  $Mn^{III}O_2Mn^{IV}]^{3+}$  reaction systems. Most reports on the oxidation of  $SCN^-$  indicate that 1 mole of  $SCN^-$  reacts with 6 moles of the oxidant yielding the reduced form of the oxidant,  $CN^-$  and  $SO_4^{2-}$  as the redox products (Stanbury et. al., 1980; Chung and Wood, 1970; Ng and Henry, 1975; Nord et. al., 1978). The oxidation of  $SCN^-$  to  $CN^-$  is suggested to pass through the intermediate species  $(SCN)_2$  and  $(SCN)_3^-$  (Ng and Henry, 1975; Barbosa-Filbo and Monhemius, 1994). Thiocyanogen  $(SCN)_2$ , is known to undergo hydrolysis in aqueous solution (Ng and Henry, 1975; Nord et. al., 1978; Olatunji

and Ayoko, 1984), as in equation (2), so that CN and  $\text{SO}_4^{2-}$  are some of the reaction products.



CN was identified by precipitating with  $\text{AgNO}_3$  while  $\text{SO}_4^{2-}$  was identified by the production of a white precipitate on addition of  $\text{BaCl}_2$  to the reacted mixture.  $\text{Mn}^{2+}$  was detected by reaction with indole (Lohdip et. al., 1998; Arabel et. al., 1997).

### Order of reaction

Kinetic measurements were made under pseudo-first-order conditions with  $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}] = 3.58 \times 10^{-4} \text{ mol dm}^{-3}$ ,  $[\text{SCN}] = (2.76 - 22.05) \times 10^{-3} \text{ mol dm}^{-3}$ ,  $[\text{H}^+] = 0.05 \text{ mol dm}^{-3}$  and  $I = 0.50$




Fig. 1: Plot of  $k_2$ , ( $\text{dm}^3 \text{mol}^{-1} \text{s}^{-1}$ ) Versus  $[\text{H}^+]$ , (mol) for the oxidation of thiocyanate ion by  $\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}$ , (a) = experimental points, (b) = regressed line.

$[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}] = 3.58 \times 10^{-4} \text{ mol dm}^{-3}$ ,  $[\text{SCN}] = 1.01 \times 10^{-2} \text{ mol dm}^{-3}$ ,  
 $I = 0.50 \text{ mol dm}^{-3}$ ,  $\lambda = 555 \text{ nm}$ ,  $T = 29.0 \pm 0.1^\circ \text{C}$ .

Table 2: Temperature dependent rate constants and activation parameters for the oxidation of thiocyanate ion by  $\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}$

T, K	293 K	297 K	302 K	307	317
$10^4 k_0, \text{s}^{-1}$	7.14	9.48	12.22	16.63	18.54

\* $E_a = 36.62 \text{ kJ mol}^{-1}$ , \* $\Delta H^\ddagger = 31.11 \text{ kJ mol}^{-1}$ , \* $\Delta S^\ddagger = -380.33 \text{ J mol}^{-1} \text{K}^{-1}$

\*values calculated at 302 K

$[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}] = 3.58 \times 10^{-4} \text{ mol dm}^{-3}$ ,  $[\text{SCN}] = 1.01 \times 10^{-2} \text{ mol dm}^{-3}$ ,  
 $[\text{H}^+] = 0.05 \text{ mol dm}^{-3}$ ,  $I = 0.50 \text{ mol dm}^{-3}$ ,  $\lambda = 555 \text{ nm}$ .

mol dm<sup>-3</sup> (NaCl). Pseudo-first order plots of log (A<sub>0</sub> - A<sub>t</sub>) versus time (where A<sub>0</sub> and A<sub>t</sub> are absorbances at time zero and t respectively), were linear to more than 80% of the extent of reaction, indicating that the reaction is first order in [SCN<sup>-</sup>]. The slope (1.07) of log-log plot of k<sub>obs</sub> versus [SCN<sup>-</sup>] supports first order in [SCN<sup>-</sup>]. The values of the second order rate constants calculated from k<sub>2</sub> = k<sub>obs</sub>/[SCN<sup>-</sup>], were found to be fairly constant (Table 1), indicating that the reaction is also first order in [Mn<sup>III</sup>O<sub>2</sub>Mn<sup>IV</sup>], giving a second order reaction overall. The rate equation for the title reaction can therefore be represented by equation (4), with k<sub>2</sub> = (1.19 ± 0.05) × 10<sup>1</sup> mol<sup>-1</sup> dm<sup>3</sup> s<sup>-1</sup> at 29.0°C.

$$-3 \frac{d[\text{oxidant}]}{dt} = k_2 [\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}] [\text{SCN}^-] \quad \dots(3)$$

Second order kinetics have also been observed for other oxidations of SCN<sup>-</sup> (Olatunji and Ayoko, 1984; Peloso et. al, 1967a; Peloso et. al., 1967b; Davies and Watkins, 1970).

The effect of acid on the rate of reaction was investigated by varying the concentration of HCl (0.01 - 0.15 mol dm<sup>-3</sup>) at [Mn<sup>III</sup>O<sub>2</sub>Mn<sup>IV</sup>] = 3.58 × 10<sup>-4</sup> mol dm<sup>-3</sup>, [SCN<sup>-</sup>] = 1.01 × 10<sup>-2</sup> mol dm<sup>-3</sup> and I = 0.50 mol dm<sup>-3</sup> (NaCl). The results (Table 1.) suggest that the rate constant increases with increase in [H<sup>+</sup>]. A least squares plot of k<sub>2</sub> versus [H<sup>+</sup>] (Fig.1) was linear (r = 0.99) with a positive intercept, so that the relationship can be given by equation (5), with a = 5.92 × 10<sup>7</sup> mol<sup>-1</sup> dm<sup>3</sup> s<sup>-1</sup> and b = 1.36 s<sup>-1</sup> at 29.0°C.

$$k_2 = a + b [\text{H}^+] \quad \dots (4)$$

The observed acid catalysis may be due to the successive protonation/hydration of the dimer (Chaudhuri et. al., 1995), since the rates of most of the oxidations of SCN<sup>-</sup> have been reported to have [OH<sup>-</sup>] dependence (Gowda and Bhat, 1988), inverse dependence on [H<sup>+</sup>] (Ng and Henry, 1975; Davies and Watkins, 1970; Kazakov and Konovalova, 1968), or independence of acid concentration (Olatunji and Ayoko, 1984; Przystas and Sutin, 1973).

Changes in ionic strength and dielectric constant of the medium had no effect on the rate of the reaction. It is expected that the rate should vary with changes in these two parameters since the redox species are charged. The non-dependence of ionic strength and dielectric constant may only be explained in terms of the Proton Coupled Electron Transfer (PCET) process in which 3 species are simultaneously reacting in the rate determining step. The addition of HCOO<sup>-</sup> and NO<sub>3</sub><sup>-</sup> also had no effect on the rate of the reaction.





The addition of a solution of acrylamide to a partially reacted mixture did not give any gel in the presence of excess methanol, indicating the probable absence of free radicals in the reaction. The absence of free radicals has also been observed for the oxidation of other sulphur containing species like methionine (Lohdip and Iyun, 1998; Ng and Henry, 1975; Lawal, 1997) and 2-mercaptobenzoic acid (Lohdip and Iyun, 2001).

The results of the spectroscopic studies indicate no significant shifts from the characteristic absorption bands of Mn<sup>III</sup>O<sub>2</sub>Mn<sup>IV</sup> (525, 555 and 648 nm). This suggests the absence of the formation of an intermediate complex in the reaction or that if such intermediate is formed, it has a small formation constant. This observation has also been reported for Mn<sup>III</sup>O<sub>2</sub>Mn<sup>IV</sup>/H<sub>2</sub>C<sub>2</sub>O<sub>4</sub> (Lohdip et. al., 1998), Mn<sup>III</sup>O<sub>2</sub>Mn<sup>IV</sup>/methionine (Lohdip and Iyun, 1998) and Mn<sup>III</sup>O<sub>2</sub>Mn<sup>IV</sup>/2-mercaptobenzoic acid (Lohdip and Iyun, 2001) systems.

Rate constants were obtained at various temperatures from which plots log k<sub>2</sub> versus 1/T and log (k<sub>2</sub>/T) versus 1/T were drawn (Fig. 2). From linear least squares analysis of the plots, the activation parameters were calculated as E<sub>a</sub> = 36.62 kJ mol<sup>-1</sup>, ΔH<sup>‡</sup> = 34.11 kJ mol<sup>-1</sup> and ΔS<sup>‡</sup> = -380.23 J K<sup>-1</sup> mol<sup>-1</sup> at 29°C. The temperature-dependent rate constants and the activation parameters are shown on Table 2.

### Mechanism

As noted earlier, the observed acid catalysis in the reaction is due to the successive protonation/hydration of the dimer, Mn<sup>III</sup>O<sub>2</sub>Mn<sup>IV</sup> (Chaudhuri et. al., 1995). Under the acid condition used for the stoichiometric and kinetic studies, the following scheme 1 is proposed for the reaction.



Scheme 1.




Fig. 2: Plot ln k<sub>0</sub> Vs 1/T & ln(k<sub>0</sub>/T) Vs 1/T for the oxidation of thiocyanate ion by Mn<sup>III</sup>O<sub>2</sub>Mn<sup>IV</sup>.  
 (a) = exptal. points, (b) = regressed line. (c) = exptal. points. (d) = regressed line.

[Mn<sup>III</sup>O<sub>2</sub>Mn<sup>IV</sup>] = 3.58 x 10<sup>-4</sup> mol dm<sup>-3</sup>, [SCN<sup>-</sup>] = 1.01 x 10<sup>-2</sup> mol dm<sup>-3</sup>,  
 [H<sup>+</sup>] = 0.05 mol dm<sup>-3</sup>, I = 0.50 mol dm<sup>-3</sup>, λ = 555 nm

Scheme 1 leads to equation (9)

$$\frac{d[\text{oxidant}]}{dt} = \{k_6 + k_7 K_5 [H^+]\} [\text{oxidant}] [\text{reductant}] \dots\dots (9)$$

$$\frac{d[\text{oxidant}]}{dt} = \{k_6 + k_7 K_5 [H^+]\}$$

Equation (9) is comparable to equation (3) with  $k_6 = a$  and  $k_7K_5 = b$ .

The absence of spectrophotometric evidence for the formation of an intermediate suggests that a precursor complex is not formed prior to electron transfer and that the redox reaction most probably occurs by the outer sphere path. The Michaelis-Menten plot of  $1/k_{obs}$  versus  $1/[RSH]$  was linear with an insignificant positive intercept, indicating the absence of a pre-electron transfer step. This absence of kinetic evidence for the formation of a precursor complex is also in favour of the outer sphere mechanism.

The absence of anion catalysis is not unexpected. As suggested earlier (Lohdip and Iyun, 1998, 2001), the simultaneous involvement of 3 species in a proton coupled electron transfer (PCET) process is subject to restrictions. This effect would not encourage the participation of any added ion in the reaction so that the presence of  $HCOO^-$  or  $NO_3^-$  had no effect on the rate of the reaction.

The absence of induced polymerization when acrylamide was added to the reaction mixture is in support of a 2 electron ( $2e^-$ ) redox process. We therefore propose that the oxidation proceeds through a  $2H^+/2e^-$  path as indicated earlier (Lohdip and Iyun, 1998, 2001; Arabel et. al., 1997; Lawal, 1997). In addition,  $[Mn^{III}O_2Mn^{IV}]$  contains a protonable moiety, a necessary condition for the occurrence of PCET.

In line with arguments presented earlier (Reddy et.al., 1994; Lohdip et. al., 1998; Lohdip and Iyun, 1998, 2001; Arabel et. al., 1997; Lawal, 1997), and those discussed above, we believe that the proton coupled electron transfer (PCET) pathway is the most probable mechanism for the reaction under investigation.

## ACKNOWLEDGMENT

We are grateful to the authorities of the University of Jos for the Grant (UJ/FNS/RG/95-96/05) provided for this research work. We are also grateful to the Ahmadu Bello University for providing facilities for this work and the University of Jos, for award of study fellowship to one of the authors (YNL).

## REFERENCES

- Arabel, J. M., Ayoko, G. A. and Singh, K., 1997. Kinetics of the Oxidation of  $\alpha$ - Hydroxyacids by High Valent Oxo-Manganese Complexes. (Unpublished).
- Arulsamy, N., Glerup, J, Hazell, A, Hodgson, D. J., McKenzie, C. J. and Toftlund, H., 1994. Synthesis and Characterization of Complexes Containing the ( $\mu$ -oxo)( $\mu$ - acetato) dimanganese (III/III) Core. *Inorg. Chem.*, 33(14): 3023 - 3025.
- Babcock, G. T., 1987. *New Comprehensive Biochemistry*, Photosynthesis, J. Amesz, Ed., Elsevier, Amsterdam pp 235 - 158.
- Barbosa-Filbo, O. and Monhemius, A. J., 1994. Leaching of Gold in Thiocyanate Solutions. Part I: Chemistry and Thermodynamics. *Trans. Inst. Min. Metal., Sect. C*, -103: C103 - C110.
- Bassett, J., Denney, R. C., Jeffrey, G. H. and Mendham, J., 1983. *Vogel's Textbook of Quantitative Inorganic Analysis: including Elementary Instrumental Analysis*, 4th Ed, The E.L.B.S. and Longman, London.
- Chaudhuri, S, Mukhopadhyay, S and Banerjee, R., 1995. Solution Equilibria and Redox Reactivities of Dioxo-bridged Manganese Complex. *J. Chem. Soc. Dalton. Trans.*, 621 - 624.
- Chung, J. and Wood, J. L., 1970. Oxidation of thiocyanate to cyanide and sulfate by the Lactoperoxidase-hydrogen peroxide system. *Arch. Biochem. Biophys.*, 142(1): 73 - 78.
- Cooper, S. R and Calvin, M., 1977. Mixed Valence Interactions in Di-oxoBridged Manganese Complexes. *J. Am. Chem. Soc.*, 99(20): 6623 - 6630.
- Dave, B. C and Czernuzewicz, R. S., 1994. Coordination Chemistry of Manganese with 2,2'-Bipyridine: Synthesis Access to High-Valent Polynuclear Oxomanganese Complexes of Biological Relevance. *New J. Chem.*, 18(1): 149 - 155.

- Davies, G and Watkins, K. O., 1970. Kinetics of Some Oxidation-Reduction Reactions Involving Cobalt(III) in Aqueous Perchloric acid. *J. Phys. Chem.*, 74: 3388 - 3392.
- Goodson, P. A, Hodgson, D. J, Glerup, J., Michelsen, K and Weihe, H., 1992. Synthesis and Characterization of Binuclear Manganese(III,IV) and Manganese(IV,IV) with 1,1,7,10-Tetra-azacyclododecane(cyclen). *Inorg. Chim. Acta.*, 197(2): 141 - 147.
- Gowda, B. T. and Bhat, J. I., 1988. Mechanism of Oxidation of Thiocyanate Ion by Bromamine-T and Hypobromite Ion in Alkaline Medium. A Kinetic Study. *J. Indian. Chem. Soc.*, 63: 159 - 163.
- Iyun, J. F, Ayoko, G. A and Lohdip, Y. N., 1997. Exploratory Kinetic Studies of the Reduction of High-valent Oxo Manganese Complexes as Model for Multinuclear Redox Active Manganese Enzymes. Paper presented at the 20th Annual Conference of the Chemical society of Nigeria. 22nd - 26th September, 1997.
- Jeffery, J. C, Thornton, P and Ward, M. D., 1994. An Unusual Chainlike Tetranuclear Manganese(II) Complexes Displaying Ferromagnetic Exchange. *Inorg. Chem.*, 33 (16): 3612.
- Kazakov, V. P. and Konovalova, U. V., 1968. Kinetics of the Reduction of Tetrabromaurate by Sulfite and Thiocyanate Ions. *Zh. Neorg. Khim.*, 13: 447 - 453.
- Lawal, H. M., 1997. The Electron Transfer Reactions of Pararosaniline Chloride (Basic Fuchsin) and Trisoxalatocobaltate(III) Ion in Acidic Media. Ph.D. Thesis, Abubakar Tafawa Balewa University, Bauchi, Nigeria.
- Lohdip, Y. N, Davies, A. K, Openshaw, J. D and Iyun, J. F., 1997. Construction and Evaluation of Solid State Photometer as an Alternative to the Conventional Spectrophotometers. *Nigerian. J. Chem. Res.*, 2, 27 - 31.
- Lohdip, Y. N and Iyun, J. F., 1998. Kinetics of the Reduction of Di- $\mu$ -tetrakis(1,10-phenanthroline)-dimanganese(III,IV) Perchlorate by DL-Methionine in Acid Solution. *Bull. Chem. Soc. Ethiop.* 12: 113 - 119.
- Lohdip, Y. N. and Iyun, J. F., 2001. Kinetics of the Oxidation of 2-Mercaptobenzoic Acid by Di- $\mu$ -oxo-tetrakis(1,10-phenanthroline)-dimanganese (III,IV) Perchlorate in Acid Medium". *Global J. Pure & Applied Sciences.*, 7(2), 291 - 196.
- Lohdip, Y. N, Iyun, J. F and Ayoko, G. A., 1998. Kinetics of the Oxidation of Oxalic Acid by Di- $\mu$ -tetrakis(1,10-phenanthroline)-dimanganese (III,IV) Perchlorate in Aqueous Acid Medium. *Indian. J. Chem.*, 37A: 1111 - 1114.
- Miao, Y. Q., Sun, X., Jiang, Z., Liao D. and Wang, G., 1995. Synthesis and Magnetism of Tetrachlorophthalato-bridged Manganese(II) Binuclear Complexes. *Transition. Met. Chem.*, 20: 270 - 275.
- Ng, F. T. T. and Henry, P. M., 1975. Kinetics and Mechanism of the Oxidation of Thiocyanate by Tris(1,10-phenanthroline)iron(III) and its Derivatives. *Can. J. Chem.*, 53: 3319 - 3326.
- Nord, G., Pedersen, B. and Farver, O., 1978. Outersphere Oxidation of Iodide and Thiocyanate by Tris(2,2-bipyridyl)- and Tris(1,10-phenanthroline)osmium(III) in Aqueous Solutions. *Inorg. Chem.*, 17: 2233 - 2238.
- Olatunji, M. A. and Ayoko, G. A. 1984. The Kinetics of the Oxidation of Iodide and Thiocyanate Ions by 12-Tungstocobaltate(III). *Polyhedron.*, 3:191 - 197.
- Peloso, A., Dolcetti, G. and Ettore, R., 1967a. Redox Reaction Mechanisms of Platinum(IV) Complexes. *Inorg. Chim. Acta.* 1, 307 - 310.
- Peloso, A., Dolcetti, G. and Ettore, R., 1967b. Reduction Reactions of Trans-PtCl<sub>2</sub>X<sub>2</sub> Complexes: Influence of the Uncharged Ligands L on the Reduction Reaction Rates. *Inorg. Chim. Acta.* 1: 403 - 406.
- Przystas, T. J. and Sutin, N., 1973. Kinetic Studies of Anion-assisted Outer-sphere Electron Transfer Reactions. *J. Amer. Chem. Soc.*, 95: 5545 - 5555.
- Reddy, K. R., Rajasekharan, V. M, Padhye, S. Dahan, F and Tuchagues, J. P., 1994. Modelling the Photosynthetic Water Oxidation Center: Synthesis, Structure and Magnetic Properties of [Mn<sub>2</sub>( $\mu$ -O)<sub>2</sub>( $\mu$ -OAc)(H<sub>2</sub>O)<sub>2</sub>(bipy)](ClO<sub>4</sub>)<sub>2</sub>.H<sub>2</sub>O. *Inorg. Chem.*, 33(3): 428 - 433.
- Stanbury, D. M., Wilmarth, W. K., Khalaf, S., Po, H. N. and Byrd, J. E., 1980. Oxidation of Thiocyanate and Iodide by Iridium (IV). *Inorg. Chem.* 19: 1715 - 2722.

- Tanase, T and Lippard, S., 1995. Dinuclear Manganese(II) Complexes with  $[\text{Mn}_2(\mu\text{-carboxylato})_2]^{2+}$  Core and Their Transformation to  $(\mu\text{-Oxo})\text{bis}(\mu\text{-carboxylato})\text{-dimanganese(III)}$  Complexes. *J. Inorg. Chem.*, 34(18): 4682.
- Projanowicz, M, Worsfold, P. J and Clinch, J. R., 1988. Solid-State Photometric Detectors for Flow Injection Analysis. *Trends. Anal. Chem.*, 7(8): 302.
- Vincent, J.B, Tsai, H. L, Blackman, A. G, Wang, S. Y, Boyd, P. W. D, Foltz, K, Huffman, J. C, Lobkovky, E. B, Hendrickson, D. N. and Christou, G., 1993. Models of the Manganese Catalase Enzymes: Dinuclear Manganese(III) Complexes with the  $[\text{Mn}_2(\mu\text{-O})(\mu\text{-O}_2\text{CR})_2]^{2+}$  Core and Terminal Monodentate Ligands: Preparation and Properties of  $[\text{Mn}_2(\mu\text{-O}_2\text{CR})_2\text{X}_2(\text{bipy})_2]$  (X = Cl,  $\text{N}_3^-$ ,  $\text{H}_2\text{O}$ ). *J. Am. Chem. Soc.*, 115(26): 12353.