

KINETICS AND MECHANISM OF THE OXIDATION OF ACETALDEHYDE BY BROMATE ION IN AQUEOUS PERCHLORIC ACID

By Y.N. Lakshmi* & A. K. Davies, Department of Chemistry & Applied Chemistry, University of Salford, M5 4WT, England.
and J.F. Iyem, Department of Chemistry, Ahmadu Bello University, Zaria, Nigeria.

ABSTRACT

The stoichiometry, kinetics and mechanism of the oxidation of acetaldehyde by bromate ion in aqueous perchloric acid have been investigated. The reaction follows the empirical rate law;

$$-d(\text{CH}_3\text{CHO})/dt = \{a + b[\text{H}^+]\} [\text{CH}_3\text{CHO}] [\text{BrO}_3^-]$$

Where $a = 2.14 \times 10^3 \text{M}^{-1} \text{s}^{-1}$, $b = 2.71 \times 10^4 \text{M}^{-2} \text{s}^{-1}$ at $23.0 \pm 0.1^\circ\text{C}$, $[\text{H}^+] = 0.05\text{--}0.15\text{M}$ and $I = 1.0\text{M}(\text{NaClO}_4)$. The mechanism of the reaction is discussed in terms of the outer-sphere mechanism.

INTRODUCTION

Several studies into the redox reactions involving metal ion complexes and a wide range of substrates have been reported¹⁻¹¹. Our interest has mostly been focused on the reactions involving oxyanions and metal ion complexes^{1-2,6}. The reactions of oxyanions with some aldehydes and ketones have also started receiving some attention¹²⁻¹⁵. In order to gain more information into their chemical and biochemical activities, we have initiated kinetic studies into the reactions of aliphatic aldehydes and ketones with selected oxyanions^{16,17}. In this paper, we report the results of the reaction between CH_3CHO and the versatile oxidizing agent, bromate ion^{3,13,18,19,21}.

EXPERIMENTAL

Reagents

Standard solutions of acetaldehyde (Analar) were usually prepared before use. Sodium bromate (Analar) solution was prepared and standardized according to the method described by Birk and Kozub^{18,19}. HClO_4 was used as the acid medium and NaClO_4 was used to maintain a constant ionic strength.

Stoichiometry

The stoichiometry of the reaction was determined by

* Author for correspondence.

† Present Address: Department of Chemistry, University of Jos, Jos, Nigeria.

spectrophotometric titration, using the mole ratio method. Reaction mixtures containing CH_3CHO and BrO_3^- solutions at $[\text{H}^+] = 0.1\text{M}$ and $I = 1.0\text{M}$ (NaClO_4), were allowed to stand until the reaction had gone to completion. The absorbances of the solutions were measured at 390nm and the stoichiometry was evaluated from the plot of the absorbance versus mole ratio.


Kinetic measurements

All rate measurements were made using Unicam SP 1750 uv-visible spectrophotometer. The reaction rates were monitored under pseudo-first order conditions, by following the rate of change of absorbance of the reaction mixture at 390nm. The pseudo-first order rate constants were obtained from the slope of the plots of $\log(A - A_\infty)$ versus time (where A_t and A_∞ are the absorbances of the reaction mixtures at time t and time zero respectively). The temperature was kept constant at $23.0 \pm 0.1^\circ\text{C}$ for all the runs, $[\text{H}^+] = 0.1\text{M}$, $I = 1.0\text{M}$ (NaClO_4), unless otherwise stated.

RESULTS AND DISCUSSION

Stoichiometry

The stoichiometry of the reaction between acetaldehyde and bromate ion was found to be in the ratio 5:2, so that the overall equation can be written as;



A similar stoichiometry was reported for the reaction between HCHO and MnO_4^- and for that between HCHO and BrO_3^- . The products of the reaction were qualitatively identified^{1,2,17}. The ether extract of the reaction mixture gave an effervescence in the presence of $NaHCO_3$. The evolved gas turned lime water milky, suggesting CO_2 , which confirmed the presence of a carboxylic acid. The brownish yellow colour of the reaction mixture was decolorized in the presence of cyclohexene, also suggesting Br_2 as one of the reaction products.

Table 1: Second order rate constants for the oxidation of acetaldehyde by bromate ion in aqueous perchloric acid at $23.0 \pm 0.1^\circ C$ $[BrO_3^-] = 0.42M, \lambda_{max} = 390nm$.

$10^2[CH_3CHO]$, (M)	$[H^+]$, (M)	I, (M)	10^2k_2 , ($M^{-1}s^{-1}$)
0.8	0.1	1.0	5.73
1.0	0.1	1.0	5.65
1.2	0.1	1.0	5.77
1.5*	0.1	1.0	5.57
1.7	0.1	1.0	5.55
1.9	0.1	1.0	5.51
2.5	0.1	1.0	5.78
1.5	0.05	1.0	3.47
1.5	0.15	1.0	6.40
1.5	0.2	1.0	7.13
1.5	0.3	1.0	11.00
1.5	0.1	0.4	4.07
1.5	0.1	0.52	4.89
1.5	0.1	0.75	5.02
1.5	0.1	1.0	5.10
1.5	0.1	1.25	5.37
1.5	0.1	1.5	5.77
1.5	0.1	2.0	5.99
1.5	0.1	2.2	6.77

Rate law

Pseudo-first order rate constants, k_0 , show a linear dependence on $[CH_3CHO]$ at constant $[H^+]$, from which the second order rate constants, k_2 , were evaluated (Table 1). The linearity of the pseudo-order rate plots and the plot of $\log k_0$ versus $\log[CH_3CHO]$, indicates that the reaction, under our

experimental conditions, is first order each in $[CH_3CHO]$ and $[BrO_3^-]$. Thus, the rate equation for the reaction is

$$-d[CH_3CHO]/dt = k_2[CH_3CHO][BrO_3^-] \dots\dots\dots(2)$$

$$k_2 = (5.65 + 0.11) \times 10^2 M^{-1} s^{-1}$$

Effect of acid

The effect of acid on the rate of the reaction was determined by keeping all other conditions constant while varying the acid concentration in the range 0.05 - 0.15M. A plot of k_2 versus $[H^+]$ was linear ($r = 0.98$), with a positive intercept, suggesting that k_2 vary with $[H^+]$ according to equation (3)

$$k_2 = a + b[H^+] \dots\dots\dots(3)$$

$$a = 2.14 \times 10^2 M^{-1} s^{-1}, b = 2.71 \times 10^3 M^{-2} s^{-1}$$

When the expression for k_2 in equation (3) is substituted into equation (2), the rate equation for the reaction becomes

$$-d[CH_3CHO]/dt = \{a + b[H^+]\} [CH_3CHO] [BrO_3^-] \dots\dots\dots(4)$$

Effect of added anions

The effect of added formate and nitrate ions on the rate of the reaction was investigated and the results, as presented in Table 2, show that both anions enhanced the rate. The enhancement of the rate is in the order $k_0(HCOO^-) > k_0(NO_3^-)$, similar to that observed for the HCHO/ BrO_3^- system¹⁷. Least squares plots of k_0 versus $[X^-]$, ($X^- = HCOO^-$ or NO_3^-), were linear and fitted into equation (5).

$$k_0 = p + q[X^-] \dots\dots\dots(5)$$

where $p(HCOO^-) = 7.27 \times 10^4 s^{-1}$,
 $q(HCOO^-) = 8.54 \times 10^2 M^{-2} s^{-1}$
 $p(NO_3^-) = 7.11 \times 10^4 s^{-1}$,
 $q(NO_3^-) = 2.28 \times 10^2 M^{-2} s^{-1}$

Table 2: Anion dependent rate constant for the oxidation of acetaldehyde by bromate ion in aqueous perchloric acid
 $[H^+] = 0.1M$, $[BrO_3^-] = 0.32M$, $[CH_3CHO] = 1.5 \times 10^{-2}M$,
 $I = 1.0M$ (NaClO₄), Temp = $23 \pm 0^\circ C$, $\lambda_{max} = 390nm$








$10^3 [X]$, (M)	$10^2 k_2$, (M ⁻¹ s ⁻¹)	(HCOO)	(NO ₃)
3.0	6.63		
5.0	7.85		5.67
7.0	8.75		6.30
10.0	10.50		6.47
15.0	.	.	6.93
20.0	.	.	7.59

Effect of ionic strength

The effect of ionic strength on the rate of the reaction was studied over the range 0.40 - 2.2M, using NaClO₄, while keeping all other conditions constant. The results presented in Table 1, show that the rate constants increase with increase in ionic strength, suggesting a positive Bronsted-Debye^{21,23} salt effect.

Reaction mechanism

The rates of redox reactions of many oxyanions show strong dependence on acid concentration^{1,3,24}. The $[H^+]$ terms commonly encountered in reactions of BrO_3^- ,^{18,19,21,25-27} has been attributed in the formation of $HBrO_3$, which reacts with the reductant in the rate determining step. In reaction systems^{17,26} similar to the titled reaction, the H^+ dependence has also been ascribed to the substantial and rapid hydration of the aliphatic aldehydes and ketones. From the above reasoning, the reaction scheme proposed below, accounts for the experimental results.



From the above, the rate of loss of CH_3CHO is given by equation (14)

$$-d[CH_3CHO]/dt = \{ [k_2 K_1^1 + k_4] + (k_1 K_1 K_1^1 + k_3 K_1) [H^+] \} [CH_3CHO] [BrO_3^-] \quad \dots\dots\dots(14)$$

where $k_1^1 = K_1 [H_2O]$ and K_1 has been reported as *ca* $10^{11} M^{-2}$.

Equation (14) is analogous to equation (4) and is consistent with the experimental results.

A key point of interest is whether this reaction occurs via the inner- or the outer-sphere mechanism or a combination of both. Cases have been reported^{18,21}, where the reaction of BrO_3^- has been explained in terms of parallel outer- and inner-sphere pathways, with the acid dependent path occurring via the outer-sphere mechanism. In some other cases¹⁹, the BrO_3^- reduction by Fe(III) complexes involve two parallel outer-sphere paths and that by $IrCl_6^{2-}$, involve essentially diffusion controlled outer-sphere processes. This issue was addressed as follows;

(a) The spectra of the reaction mixtures taken 2 minutes after initiation of the reaction did not show any shift in the λ_{max} suggesting that there is no inner-sphere complex formation^{1,17}. This observation indicates the occurrence of the outer-sphere mechanism

(b) Addition of acrylonitrile to the partially

oxidized reaction mixture of CH_3CHO and BrO_3^- showed no gel formation, even on addition of large excess of methanol^{29,30}. This suggests the absence of free radicals in the reaction.

(c) From Table 2, it can be observed that the rate of the reaction was enhanced by the presence of added formate and nitrate ions in the following order; $k(\text{HCOO}^-) > k(\text{NO}_3^-)$. This anion catalysis is in support of the outer-sphere mechanism^{32,31}. Similar anion catalysis has been observed for other BrO_3^- systems¹²⁷.

(d) A Lineweaver-Burk modification³² of Michaelis-Menten plot of $1/k$ versus $1/[\text{CH}_3\text{CHO}]$ was linear with an insignificant intercept. This also suggests that inner-sphere intermediates are unimportant in this reaction.

From the above reasoning (a) - (d), it can be inferred that the spectroscopic and catalytic evidence are in support of the occurrence of the outer-sphere mechanism in the titled reaction.

Acknowledgement

We are grateful to the University of Salford for providing the facilities for this work and to the University of Jos for the award of study fellowship to YNL.

REFERENCES

1. Iyun, J.F., Ayoko, G.A. and Lohdip, Y.N., *Polyhedron*, 1992, 11 (18), 2389.
2. Iyun, J.F., Ayoko, G.A. and Lawal, H.M., *Transition. Met. Chem.*, 1992, 12, 63.
3. Ayoko, G.A., Iyun, J.F. and El-Idris, I.F., *Transition. Met. Chem.*, 1991, 16, 145.
4. Ayoko, G.A., Iyun, J.F. and El-Idris, I.F., *Transition. Met. Chem.*, 1992, 17, 324.
5. Iyun, J.F., Ayoko, G.A. and Ekubo, A., *Indian J. Chem.*, 1992, 31A, 943.
6. Iyun, J.F., Ayoko, G.A. and Lawal, H.M., *Indian J. Chem.*, 1992, 31A, 975.
7. Ayoko, G.A., Iyun, J.F. and El-Idris, I.F., *Transition, Met. Chem.*, 1993, 18, 278.
8. Ayoko, G.A., Iyun, J.F. and Mamman, S., *Transition Met. Chem.*, 1993, 18, 475.
9. Iyun, J.F., Ayoko, G.A. and Lawal, H.M., *Transition Met. Chem.*, 1995, k20, 30.
10. Iyun, J.F., Lawal, H.M. and Olagbemiro, O.T., *Indian J. Chem.*, 1995, 34A, 446.
11. Iyun, J.F., Musa, K.Y. and Ayoko, G.A., *Indian J. Chem.*, 1995, 31A, 635.
12. Banerji, K.K., *Indian J. Chem.*, 1973, 11, 242.
13. Panigrahi, G. and Misro, P.K., *Indian J. Chem.*, 1977, 15A, 1066.
14. Radhakrishnamurti, P.S. and Sahu, B., *Indian J. Chem.*, 1977, 15A, 700.
15. Singh, M.P., Singh, D.R., Singh, A.K. and Kumar, M., *Indian J. Chem.*, 1977, 15A, 718.
16. Iyun, J.F. and Ajibade, F., *J. Chem. Soc. Nig.*, (1994). In Press.
17. Lohdip, Y.N., Iyun, J.F. and Mutihir, I.J., *Ghana J. Chem.*, (1995). In Press.
18. Birk, J.P. and Kozub, S.G., *Inorg. Chem.*, 1973, 12, 2489.
19. Birk, J.P. and Kozub, S.G., *Inorg. Chem.*, 1988, 27, 1186.
20. Sharp, C. and Skyes, A.G., *Inorg. Chem.*, 1978, 17, 501.
21. Birk, J.P., *Inorg. Chem.*, 1973, 12, 2486.
- 22a. Bronsted, J.N., *Z. Phys. Chem.*, k1922, 102, 160.
- 22b. Wilkins, R.G., *Kinetics and Mechanism of Reactions of Transition Metal Complexes*, 2nd Thoroughly Revised Edition, VCH, p110 - 118, 1991.
- 22c. Adegite, A., Iyun, J.F. and Ojo, J.F., *J. Chem. Soc. Dalton Trans.*, 1977, 11.
23. Birk, J.P., *Inorg. Chem.*, 1978, 17, 504.
24. Thompson, R.C., *Inorg. Chem.*, 1971, 10, 1892.
25. Knight, G.C. and Thompson, R.C., *Inorg. Chem.*, 1971, 10, 1892.
26. Goswami, C. and Banerji, K.K., *Bull. Chem. Soc. Japan*, 1973, 43, 2643.
27. Ayoko, G.A. and Olatunji, M.A., *Transition Met. Chem.*, 1985, 10, 218.
28. Sen-Gupta, K.K., Das, S. and Sen-Gupta, S., *Transition Met. Chem.*, 1978, 12, 417.
29. Przystas, T.J. and Sutin, N., *J. Amer. Chem. Soc.*, 1973, 95, 5545.
30. Henri, V., *Compt. Rend. Acad. Sci. Paris.*, 1902, 135, 916.
31. Michaelis, L. and Menten, M.L., *Biochem. Z.*, 1913, 49, 33.

accepted 19/9/96
received 29/5/96